


A Phenomenological Model for the Glasma and Photons

L. McLerran, Crete, 2014



← **Strongly Interacting QGP** →

The Glasma are highly coherent colored fields evolving to a thermalized QGP
 The Glasma is weakly coupled but strongly interacting

$$\alpha_S \ll 1 \quad A \sim \frac{1}{g} Q_{sat} \quad \frac{dN}{dyd^2r_T} \sim \frac{Q_{sat}^2}{\alpha_S}$$



RUPRECHT-KARLS-
UNIVERSITÄT
HEIDELBERG



How Does the Glasma Evolve:

At an early time:

$$\frac{1}{\tau\pi R^2} \frac{dN}{d^3p_T} = f(p)$$

$$f(p) \sim \frac{1}{\alpha_s}, \quad p \leq Q_{sat}$$

$$f(p) \leq 1, \quad p \geq Q_{sat}$$

System evolves by scattering and two scales emerge

$$\Lambda_{IR}, \quad f(\Lambda_{IR}) \sim \frac{1}{\alpha_S}$$

$$\Lambda_{UV}, \quad f(\Lambda_{UV}) \sim 1$$

How do these scales evolve?

In transport equation:

$$df/dt \sim \alpha^2 f^3$$

The term with four factors of f cancels in the difference between backwards and forward going processes

If the process is dominated in the infrared:

$$df/dt \sim \frac{1}{\tau_{scat}} f$$

The scattering time can be evaluated in terms of the two scales by explicitly evaluating the phase space integrals in the transport equations

$$\tau_{scat} \sim \frac{\Lambda_{UV}}{\Lambda_{IR}} \frac{1}{\Lambda_{IR}}$$

Note that factors of coupling strength have disappeared. The scattering time is the Lorentz time dilation of the infrared scattering scale when the coherence is maximal. This result is true also when including inelastic scattering.

The equation:

$$\tau_{scat} \sim \frac{\Lambda_{UV}}{\Lambda_{IR}} \frac{1}{\Lambda_{IR}}$$

Is true except close to a thermal fixed point. Near a thermal fixed point, the right hand side of the transport equation vanishes. Near the thermal fixed point, the evolution of the system slows as one has approached equilibrium.

Far from equilibrium, we expect

$$\tau \sim \tau_{scat}$$

We will soon see that the time evolution of both scales is determined by this condition and the condition of energy conservation, assuming that in the infrared, the distributions functions are classical thermal distribution functions

$$f \sim \frac{1}{\alpha_S} \frac{\Lambda_{IR}}{E} \quad \epsilon \sim \int d^3p p f \sim \frac{1}{\alpha_S} \Lambda_{IR} \Lambda_{UV}^3$$

A simple model, assuming local equilibration in the infrared is

$$f(p) = \frac{\kappa \Lambda_{IR}}{\alpha_S \Lambda_{UV}} \frac{1}{e^{E/\Lambda_{UV}} - 1}$$

This distribution is a classical thermal distribution in the infrared

$$f \sim \frac{1}{\alpha_S} \frac{\Lambda_{IR}}{E}$$

and goes to zero when $E \sim \Lambda_{UV}$

It is like a thermal; distribution with a temperature $T \sim \Lambda_{UV}$

It becomes a thermal distribution function when the over-occupation factor $\frac{\kappa \Lambda_{IR}}{\alpha_S \Lambda_{UV}} \rightarrow 1$

Or when $\kappa \Lambda_{IR} = \alpha_S \Lambda_{UV}$

Then the infrared scale is that of the magnetic mass and the UV scale is the temperature

Note that the entropy of the gluon distribution is

$$s \sim \int d^3p \{ (1+f) \ln(1+f) - f \ln(f) \} \sim \Lambda_{UV}^3 \ln \frac{\Lambda_{IR}}{\alpha_S \Lambda_{UV}}$$

But the number of gluons is

$$\rho \sim \frac{1}{\alpha_S} \Lambda_{IR} \Lambda_{UV}^2$$

So the entropy to particle ratio is less than one until thermalization
due to the coherence

$$s/n \sim \alpha_S \Lambda_{UV} / \Lambda_{IR}$$

For fermions we can use

$$q = \frac{1}{e^{E/\Lambda_{UV}} + 1}$$

The ratio of the number of quarks to gluons is suppressed until thermalization due to the over-occupation of gluonic states

$$q/g \sim \alpha_S \Lambda_{UV} / \Lambda_{IR}$$


The advantage of this parameterization of the gluon distribution functions is that thermal results can be reproduced simply by replacing the temperature with the ultraviolet scale, and multiplying the gluon distribution function by the over-occupation factor. An example of how this works is with photon production.

The Problem with Photons at RHIC and LHC

Decay photons (in pp and AA):

$$m \rightarrow \gamma + X, \quad m = \pi^0, \eta, \omega, \eta', a_1, \dots$$

Direct photons: (inclusive(=total) – decay) – measured experimentally



hard photons:

(large p_T ,
in pp and AA)

- **prompt** (pQCD; initial hard N+N scattering)
- **jet fragmentation** (pQCD; qq, gq bremsstrahlung)
(in AA can be modified by parton energy loss in medium)

thermal photons:

(low p_T , in AA)

- **QGP**
- **Hadron gas**

jet- γ -conversion in plasma


(large p_T , in AA)

jet-medium photons

(large p_T , in AA) - scattering of hard partons with thermalized partons





$$q_{\text{hard}} + q_{\text{QGP}} \rightarrow \gamma + q,$$

$$q_{\text{hard}} + q_{\text{bar QGP}} \rightarrow \gamma + q$$





Phys. Rev. C 81, 034911 (2010)

It is not clear whether the photons seen are emitted early or late, nor the source of these photons: misidentified hadron decays, jet fragmentation, QGP or hadron gas. The photons also have a large flow that is problematic. There are problems both with absolute rates and with the magnitude of v_2




There is geometric scaling of the p_T spectrum for pp, dAu, Au-Au at various hadron multiplicities and PbPb at LHC



$$Q_{sat}^2 = \frac{\kappa}{\pi R^2} \frac{dN}{dy} \quad \frac{1}{\pi R^2} \frac{dN_\gamma}{dy d^2 p_T} = F \left(\frac{Q_{sat}}{p_T} \right)$$


We also agree with the multiplicity dependence seen in Phenix
LDM and Chritian Klein -Boesing

With Bjoern Schenke we computed spectrum of photons in 1+1 hydro. Shape fits well, but the rate requires a large k factor of about 7




Because the Glasma decays more slowly than the thermalize QGP, we get acceptable flow from Glasma + QGP

The rate problem remains, but perhaps is solved by properly doing jet quenching plus fragmentation photons. A large uncertainty here is associated with how the jet contribution is computed.



Paquet, LDM and Schenke



Cocktail subtraction is very sensitive. Paquet argued at QM that McGill computation when following experimentalists prescription for subtracting the cocktail of hadron decay can fit the photon v_2 . Perhaps the photons do really come from early time? If so is it more jets or QGP or thermalized QGP or Glasma?

